附录1　影响黑颈鹤筑巢栖息地偏好的环境因子间相关性系数(1、2、3…代表左侧对应的环境因子)

Appendix 1　The correlation coefficient between environmental variables which influence nesting habitat preference of black-necked cranes (1, 2, 3… represent left corresponding variables)

	2	3	4	5	6	7	8	9	10	11	12	13
1 距深水沼泽距离 Distance to deep water marsh	0.71	–0.14	0.45	0.48	–0.01	0.15	–0.15	–0.12	–0.21	0.00	–0.09	–0.05
2 距浅水沼泽距离 Distance to shallow water marsh		–0.07	0.34	0.36	–0.14	0.12	–0.04	–0.08	0.19	–0.04	–0.13	–0.05
3 距盐化草甸距离 Distance to salinized meadow			0.09	0.20	0.06	–0.18	0.09	–0.01	–0.07	0.04	–0.00	–0.14
4 距沼泽化草甸距离 Distance to marsh meadow				0.61	0.09	0.13	–0.01	0.01	–0.03	0.09	–0.03	–0.03
5 距湖泊距离 Distance to lake					0.06	0.02	–0.19	–0.10	–0.10	0.45	–0.13	–0.03
6 距河流距离 Distance to river						0.17	0.08	–0.15	–0.13	0.25	–0.04	–0.03
7 距山脉距离 Distance to mountain							0.17	–0.51	–0.03	–0.01	–0.35	–0.18
8 距戈壁距离 Distance to gobi								0.06	0.27	–0.06	0.08	0.19
9 距公路距离 Distance to road									–0.03	–0.03	0.57	0.28
10 距房屋距离 Distance to house										–0.09	0.06	0.01
11 海拔 Altitude											–0.12	–0.03
12 坡度 Slope												0.54
13 坡向 Aspect												