

附录 1 荷叶铁线蕨简化基因组平均 GC 含量

Appendix 1 Average GC content of *Adiantum nelumboides* for GBS reduced-representation genome sequencing

居群 Population	采样地点 Sampling localities	样品采集号 Sample voucher no.	GC 含量 GC content (%)
石柱 SZ	重庆市石柱县西沱镇 Xituo Town, Shizhu County, Chongqing	YYH15105	40.8
		YYH15106	40.3
		YYH15107	40.6
		YYH15108	41.4
		YYH15109	40.4
		YYH15110	41.1
		YYH15111	41.3
		YYH15112	41.6
		YYH15113	40.1
YYH15116	40.5		
南川 NC	重庆市南川区 Nanchuan District, Chongqing	YSR0101	40.2
		YSR0102	39.9
		YSR0103	39.2
新乡 XX	重庆市万州区新乡镇 Xinxiang Town, Wanzhou District, Chongqing	YYH15071	40.6
		YYH15072	40.2
		YYH15086	39.9
		YYH15087	40.5
		YYH15088	40.2
		YYH15090	40.4
		YYH15091	39.3
YYH15092	40.2		
燕山 YS	重庆市万州区燕山乡 Yanshan Town, Wanzhou District, Chongqing	YYH15070	40.2
武陵 WL	重庆市万州区武陵镇 Wuling Town, Wanzhou District, Chongqing	YYH15117	40.2
		YYH15118	40.1
		YYH15119	40.9
		YYH15120	40.8
		YYH15121	40.2
忠县 ZX	重庆市忠县石宝镇 Shibao Town, Zhong County, Chongqing	YYH15114	40.6