

## 附录 1 非濒危植物种群核酸多样性的比较

Appendix 1 Comparison of nucleic acid diversity of non-threatened species groups

学名 Scientific name	中名 Chinese name	$\theta\pi$	样本量 Sample size	参考文献 References
<i>Populus tremuloides</i>	颤杨	0.016	22	<a href="#">Wang et al, 2016</a>
<i>Rhododendron delavayi</i>	马缨杜鹃	0.01297	30	<a href="#">Ma et al, 2021</a>
<i>Populus tremula</i>	欧洲山杨	0.0147	24	<a href="#">Wang et al, 2016</a>
<i>Phoenix dactylifera</i>	海枣	0.0092	62	<a href="#">Hazzouri et al, 2015</a>
<i>Pyrus pyrifolia</i>	沙梨	0.00173	312	<a href="#">Zhang et al, 2021</a>
<i>Prunus avium</i>	车厘子	0.00231	21	<a href="#">Xanthopoulou et al, 2020</a>
<i>Cannabis sativa</i>	大麻	0.00387	110	<a href="#">Ren et al, 2021</a>
<i>Populus trichocarpa</i>	毛果杨	0.0041	544	<a href="#">Evans et al, 2014</a>
<i>Amborella trichopoda</i>	无油樟	0.0021	12	<a href="#">Amborella Genome Project, 2013</a>
<i>Prunus persica</i>	桃	0.0015	11	<a href="#">International Peach Genome Initiative et al, 2013</a>
<i>Phoenix sylvestris</i>	樟子松	0.0105	7	<a href="#">Flowers et al, 2019</a>
<i>Phoenix canariensis</i>	海岸松	0.0117	6	<a href="#">Flowers et al, 2019</a>
<i>Musa itinerans</i>	阿宽蕉	0.0064	24	<a href="#">Wu et al, 2018</a>
<i>Vitis</i>	野生葡萄	0.0035	64	<a href="#">Liang et al, 2019</a>
<i>Glycine max</i>	野生大豆	0.00294	62	<a href="#">Zhou et al, 2015</a>
<i>Coffea canephora</i>	中粒咖啡	0.00362	34	<a href="#">Huang et al, 2020</a>
<i>Coffea excelsa</i>	高种咖啡	0.00287	11	<a href="#">Huang et al, 2020</a>
<i>Lactuca serriola</i>	野莴苣	0.00294	199	<a href="#">Wei et al, 2021</a>
<i>Solanum pimpinellifolium</i>	醋栗番茄	0.00323	53	<a href="#">Lin et al, 2014</a>
<i>Malus sylvestris</i>	欧洲野苹果	0.00255	10	<a href="#">Duan et al, 2017</a>
<i>Malus sieversii</i>	新疆野苹果	0.0013	29	<a href="#">Duan et al, 2017</a>
<i>Boehmeria nivea</i>	野生苎麻	0.00618	14	<a href="#">Wang et al, 2021</a>
<i>Ziziphus acidojujuba</i>	野生枣	0.00185	202	<a href="#">Guo et al, 2021</a>
<i>Citrullus colocynthis</i>	药西瓜	0.00675	15	<a href="#">Guo et al, 2019</a>
<i>Momordica charantia</i>	野生小果苦瓜	0.00659	21	<a href="#">Cui et al, 2020</a>
<i>Ricinus communis</i>	野生蓖麻	0.00195	56	<a href="#">Xu et al, 2021</a>
<i>Populus adenopoda</i>	响叶杨	0.004092675	27	<a href="#">Liu et al, 2022</a>
<i>Populus alba</i>	银白杨	0.005715963	17	<a href="#">Liu et al, 2022</a>
<i>Populus davidiana</i>	山杨	0.010984953	34	<a href="#">Liu et al, 2022</a>
<i>Populus qiongdaoensis</i>	琼岛杨	0.001389251	14	<a href="#">Liu et al, 2022</a>
<i>Populus rotundifolia</i>	清溪杨	0.008184706	41	<a href="#">Liu et al, 2022</a>

$\theta\pi$ : 群体中任意两条不同序列(个体)的碱基差异数(SNP)的平均值。

$\theta\pi$ : average number of pairwise nucleotide differences per site.

## 参考文献

Amborella Genome Project (2013) The *Amborella* genome and the evolution of flowering plants. *Science*, 342, 1241089.

Cui JJ, Yang Y, Luo SB, Wang L, Huang RK, Wen QF, Han XX, Miao NS, Cheng JW, Liu ZJ, Zhang CY, Feng CC, Zhu HS, Su JW,

景昭阳, 程可光, 舒恒, 马永鹏, 刘平丽 (2023) 全基因组重测序方法在濒危植物保护中的应用. 生物多样性, 31, 22679. <https://www.biodiversity-science.net/CN/10.17520/biods.2022679>.

Wan XJ, Hu F, Niu Y, Zheng XM, Yang YL, Shan D, Dong ZS, He WM, Dhillon NPS, Hu KL (2020) Whole-genome sequencing provides insights into the genetic diversity and domestication of bitter gourd (*Momordica* spp.). Horticulture Research, 7, 85.

Duan N, Bai Y, Sun H, Wang N, Ma Y, Li M, Wang X, Jiao C, Legall N, Mao L, Wan S, Wang K, He T, Feng S, Zhang Z, Mao Z, Shen X, Chen X, Jiang Y, Wu S, Yin C, Ge S, Yang L, Jiang S, Xu H, Liu J, Wang D, Qu C, Wang Y, Zuo W, Xiang L, Liu C, Zhang D, Gao Y, Xu Y, Xu K, Chao T, Fazio G, Shu H, Zhong GY, Cheng L, Fei Z, Chen X (2017) Genome re-sequencing reveals the history of apple and supports a two-stage model for fruit enlargement. Nature Communications, 8, 249.

Evans LM, Slavov GT, Rodgers-Melnick E, Martin J, Ranjan P, Muchero W, Brunner AM, Schackwitz W, Gunter L, Chen JG, Tuskan GA, DiFazio SP (2014) Population genomics of *Populus trichocarpa* identifies signatures of selection and adaptive trait associations. Nature Genetics, 46, 1089–1096.

Flowers JM, Hazzouri KM, Gros-Balthazard M, Mo ZY, Kouroumpa K, Perrakis A, Ferrand S, Khierallah HSM, Fuller DQ, Aberlenc F, Fournaraki C, Purugganan MD (2019) Cross-species hybridization and the origin of North African date palms. Proceedings of the National Academy of Sciences, USA, 116, 1651–1658.

Guo MX, Zhang ZR, Li SP, Lian Q, Fu PC, He YL, Qiao JX, Xu KK, Liu LP, Wu MY, Du ZR, Li SN, Wang JJ, Shao PY, Yu Q, Xu G, Li DK, Wang YK, Tian S, Zhao J, Feng X, Li RQ, Jiang WK, Zhao XS (2021) Genomic analyses of diverse wild and cultivated accessions provide insights into the evolutionary history of jujube. Plant Biotechnology Journal, 19, 517–531

Guo S, Zhao S, Sun H, Wang X, Wu S, Lin T, Ren Y, Gao L, Deng Y, Zhang J, Lu X, Zhang H, Shang J, Gong G, Wen C, He N, Tian S, Li M, Liu J, Wang Y, Zhu Y, Jarret R, Levi A, Zhang X, Huang S, Fei Z, Liu W, Xu Y (2019) Resequencing of 414 cultivated and wild watermelon accessions identifies selection for fruit quality traits. Nature Genetics, 51, 1616–1623.

Hazzouri KM, Flowers JM, Visser HJ, Khierallah HSM, Rosas U, Pham GM, Meyer RS, Johansen CK, Fresquez ZA, Masmoudi K, Haider N, El Kadri N, Idaghdour Y, Malek JA, Thirkhill D, Markhand GS, Krueger RR, Zaid A, Purugganan MD (2015) Whole genome re-sequencing of date palms yields insights into diversification of a fruit tree crop. Nature Communications, 6, 8824.

Huang LF, Wang XY, Dong YP, Long YZ, Hao CY, Yan L, Shi T (2020) Resequencing 93 accessions of coffee unveils independent and parallel selection during *Coffea* species divergence. Plant Molecular Biology, 103, 51–61.

International Peach Genome Initiative; Verde I, Abbott AG, Scalabrin S, Jung S, Shu SQ, Marroni F, Zhebentyayeva T, Dettori MT, Grimwood J, Cattonaro F, Zuccolo A, Rossini L, Jenkins J, Vendramin E, Meisel LA, Decroocq V, Sosinski B, Prochnik S, Mitros T, Policriti A, Cipriani G, Dondini L, Ficklin S, Goodstein DM, Xuan PF, Del Fabbro C, Aramini V, Copetti D, Gonzalez S, Horner DS, Falchi R, Lucas S, Mica E, Maldonado J, Lazzari B, Bielenberg D, Pirona R, Miculan M, Barakat A, Testolin R, Stella A, Tartarini S, Tonutti P, Arús P, Orellana A, Wells C, Main D, Vizzotto G, Silva H, Salamini F, Schmutz J, Morgante M, Rokhsar DS

景昭阳, 程可光, 舒恒, 马永鹏, 刘平丽 (2023) 全基因组重测序方法在濒危植物保护中的应用. 生物多样性, 31, 22679. <https://www.biodiversity-science.net/CN/10.17520/biods.2022679>.

(2013) The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. *Nature Genetics*, 45, 487–494.

Liang ZC, Duan SC, Sheng J, Zhu SS, Ni XM, Shao JH, Liu CH, Nick P, Du F, Fan PG, Mao RZ, Zhu YF, Deng WP, Yang M, Huang HC, Liu YX, Ding YQ, Liu XJ, Jiang JF, Zhu YY, Li SH, He XH, Chen W, Dong Y (2019) Whole-genome resequencing of 472 *Vitis* accessions for grapevine diversity and demographic history analyses. *Nature Communications*, 10, 1190.

Lin T, Zhu GT, Zhang JH, Xu XY, Yu QH, Zheng Z, Zhang ZH, Lun YY, Li S, Wang XX, Huang ZJ, Li JM, Zhang CZ, Wang TT, Zhang YY, Wang AX, Zhang YC, Lin K, Li CY, Xiong GS, Xue YB, Mazzucato A, Causse M, Fei ZJ, Giovannoni JJ, Chetelat RT, Zamir D, Städler T, Li JF, Ye ZB, Du YC, Huang SW (2014) Genomic analyses provide insights into the history of tomato breeding. *Nature Genetics*, 46, 1220–1226.

Liu SY, Zhang L, Sang YP, Lai Q, Zhang XX, Jia CF, Long ZQ, Wu JL, Ma T, Mao KS, Street NR, Ingvarsson PK, Liu JQ, Wang J (2022) Demographic history and natural selection shape patterns of deleterious mutation load and barriers to introgression across *Populus* genome. *Molecular Biology and Evolution*, 39, msac008.

Ma H, Liu Y, Liu D, Sun W, Liu X, Wan Y, Zhang X, Zhang R, Yun Q, Wang J, Li Z, Ma Y (2021) Chromosome-level genome assembly and population genetic analysis of a critically endangered rhododendron provide insights into its conservation. *Plant Journal*, 107, 1533–1545.

Ren GP, Zhang X, Li Y, Ridout K, Serrano-Serrano ML, Yang YZ, Liu A, Ravikanth G, Ali Nawaz M, Mumtaz AS, Salamin N, Fumagalli L (2021) Large-scale whole-genome resequencing unravels the domestication history of *Cannabis sativa*. *Science Advances*, 7, eabg2286.

Wang J, Street NR, Scofield DG, Ingvarsson PK (2016) Natural selection and recombination rate variation shape nucleotide polymorphism across the genomes of three related *Populus* species. *Genetics*, 202, 1185–200.

Wang YZ, Li F, He QY, Bao ZG, Zeng Z, An D, Zhang T, Yan L, Wang HY, Zhu SY, Liu TM (2021) Genomic analyses provide comprehensive insights into the domestication of bast fiber crop ramie (*Boehmeria nivea*). *The Plant Journal*, 107, 787–800.

Wei T, van Treuren R, Liu XJ, Zhang ZW, Chen JJ, Liu Y, Dong SS, Sun PN, Yang T, Lan TM, Wang XG, Xiong ZQ, Liu YQ, Wei JP, Lu HR, Han SP, Chen JC, Ni XM, Wang J, Yang HM, Xu X, Kuang HH, van Hintum T, Liu X, Liu H (2021) Whole-genome resequencing of 445 *Lactuca* accessions reveals the domestication history of cultivated lettuce. *Nature Genetics*, 53, 752–760.

Wu W, Ng WL, Yang JX, Li WM, Ge XJ (2018) High cryptic species diversity is revealed by genome-wide polymorphisms in a wild relative of banana, *Musa itinerans*, and implications for its conservation in subtropical China. *BMC Plant Biology*, 18, 194.

Xanthopoulou A, Manioudaki M, Bazakos C, Kissoudis C, Farsakoglou AM, Karagiannis E, Michailidis M, Polychroniadou C,

景昭阳, 程可光, 舒恒, 马永鹏, 刘平丽 (2023) 全基因组重测序方法在濒危植物保护中的应用. 生物多样性, 31, 22679. <https://www.biodiversity-science.net/CN/10.17520/biods.2022679>.

- Zambounis A, Kazantzis K, Tsaftaris A, Madesis P, Aravanopoulos F, Molassiotis A, Ganopoulos I (2020) Whole genome resequencing of sweet cherry (*Prunus avium* L.) yields insights into genomic diversity of a fruit species. Horticulture Research, 7, 60.
- Xu W, Wu D, Yang TQ, Sun C, Wang ZQ, Han B, Wu SB, Yu AM, Chapman MA, Muraguri S, Tan Q, Wang WB, Bao ZG, Liu AZ, Li DZ (2021) Genomic insights into the origin, domestication and genetic basis of agronomic traits of castor bean. Genome Biology, 22, 113.
- Zhang MY, Xue C, Hu HJ, Li JM, Xue YS, Wang RZ, Fan J, Zou C, Tao ST, Qin MF, Bai B, Li XL, Gu C, Wu S, Chen X, Yang GY, Liu YY, Sun MY, Fei ZJ, Zhang SL, Wu J (2021) Genome-wide association studies provide insights into the genetic determination of fruit traits of pear. Nature Communications, 12, 1144.
- Zhou ZK, Jiang Y, Wang Z, Gou ZH, Lyu J, Li WY, Yu YJ, Shu LP, Zhao YJ, Ma YM, Fang C, Shen YT, Liu TF, Li CC, Li Q, Wu M, Wang M, Wu YS, Dong Y, Wan WT, Wang X, Ding ZL, Gao YD, Xiang H, Zhu BG, Lee SH, Wang W, Tian ZX (2015) Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. Nature Biotechnology, 33, 408–414.